



RANGE

Biology and Management of the Toadflaxes no. 3.114

by K.G. Beck¹

Quick Facts...

Yellow toadflax (*Linaria vulgaris*) and the Dalmatian toadflaxes (*Linaria dalmatica* and *Linaria genistifolia*) are invasive, perennial weeds that are noxious in Colorado and other western states.

Seeds of yellow toadflax germinate and emerge in early to mid-May while Dalmatian toadflax seeds may germinate and emerge earlier, especially on south or southeast facing slopes.

Dalmatian toadflax may be controlled with Tordon 22K at 2 pt/A sprayed at flowering or in fall.

Yellow toadflax (*Linaria vulgaris*) and the Dalmatian toadflaxes (*Linaria dalmatica* and *Linaria genistifolia*) are invasive, perennial weeds that are noxious in Colorado and many other western states. Toadflax invasion is favored by disturbance and they invade degraded areas such as roadsides, abandoned lots and fields, gravel pits, clearings, and overgrazed rangeland. In Colorado, these weed species are found at elevations from 5,000 feet to over 10,000 feet. Yellow toadflax in particular has spread into high mountain valleys and parks. Yellow toadflax infests 40,800 acres in Colorado and Dalmatian toadflax infests 34,200 acres.

Origin and History

The toadflaxes have a storied past and a long relationship with humans. Dalmatian toadflax is native to the Mediterranean region. Broad-leaved Dalmatian toadflax (*L. dalmatica*) has been cultivated as an ornamental for at least 400 years. It was introduced into the western U.S. as an ornamental in 1874. The majority of Dalmatian toadflax infestations in the west are broad-leaved Dalmatian toadflax; however in its native Eurasian habitat, narrow-leaved Dalmatian toadflax (*L. genistifolia*) is more widespread, indicating




Figure 1. YELLOW TOADFLAX, butter-and-eggs, wild snapdragon (*Linaria vulgaris* Mill.). Noxious. Figwort family. This introduced creeping perennial is an escaped ornamental that reproduces by seed and extensive horizontal roots. Stems are smooth, erect, leafy, often in clumps, and 1 to 2 1/2 feet tall. Numerous pale green leaves are alternate, narrow, pointed at both ends, and 2 1/2 or more inches long. Flowers resemble those of cultivated snapdragon; each has a spur extending below from the lower lip of the corolla. They are about 1 inch long, bright yellow with a bearded, orange throat and occur in terminal, somewhat elongated clusters with the youngest flowers at the tip. The fruit is a brown, globe-shaped, two-celled capsule, 1/4 inch in diameter containing many seeds. Seeds are small, round, rough, flattened, with a papery, notched circular wing, dark brown, and about 1/12 inch in diameter. It is a persistent, aggressive invader in Colorado from 6,000 to 8,500 feet mostly on the Western Slope.

its potential to also invade and become widely dispersed in the U.S. Narrow-leaved Dalmatian toadflax currently infests several areas in western Oregon, northwestern Washington, British Columbia, and possibly in Colorado.

Yellow toadflax is native to south-central Eurasia where it was used for fabric dyes and for medicinal purposes. It was imported into North America in the late 1600s as an ornamental and for folk remedies. It was widely distributed in North America by the mid 20th century. Unfortunately in states where yellow toadflax is not noxious, it still is sold by some nurseries as “butter and eggs” or as “wild snapdragons.”

BIOLOGY

Seeds of yellow toadflax germinate and emerge in early to mid-May while Dalmatian toadflax seeds may germinate and emerge earlier especially on south or southeast facing slopes.

Germination and Emergence

Seeds of yellow toadflax germinate and emerge in early to mid-May while Dalmatian toadflax seeds may germinate and emerge earlier especially on south or southeast facing slopes. In Washington, Dalmatian toadflax seedlings on south facing slopes usually emerge in early to mid-March. First year Dalmatian toadflax plants often produce prostrate shoots in fall that survive into the following spring. Mature Dalmatian toadflax may produce prostrate shoots, but to a lesser extent, and these typically die before winter and shoots emerge the following spring from roots. Yellow and Dalmatian toadflax shoots that grow from roots emerge as early as mid-March along the Front Range in Colorado, but vegetative shoot emergence may not begin until mid- to late June at 9,000 feet to 10,000 feet. In Canada, vegetative shoots begin to emerge when soil temperatures range from 42 to 50 F.

Root growth

Seedling root develop is slow and represents a life stage vulnerable to control attempts and plant competition. Disturbance promotes toadflax invasion and may be necessary for establishment to occur. However once established, toadflaxes readily spread into adjacent non-disturbed areas. Much of this spread is by vegetative means, reflecting a vigorously-growing root system. Dalmatian toadflax roots may grow 20 inches deep or more nine weeks after seedlings have emerged and

Dalmatian toadflax typically flowers beginning in late May or June in Colorado and may continue until fall, particularly if moisture is not limiting. Yellow toadflax begins to flower when shoots are from 16 to 24 inches tall, mid- to late May along the Front Range in Colorado, although at higher elevations (9,000 feet or more), flowering may not begin until late July.



Figure 2: DALMATION TOADFLAX [*Linaria dalmatica* (L.) Mill. = *Linaria genistifolia* ssp. *dalmatica* (L.) Maire & Petitmengin]. Noxious. This differs from yellow toadflax principally in being larger and having differently shaped leaves. Clumps of stems are 3 to 4 feet tall. Waxy leaves are broad, ovate, sometimes heart shaped and upper leaves clasp the stem (an important difference). Seeds are irregular in shape, angular, somewhat flattened, thin-edged, strongly netted, tan-gray and 1/24 to 1/16 inch across. It is not as common, but is more aggressive than yellow toadflax. It is reported in various parts of the state from 5,000 to 6,500 feet.

Dalmatian toadflax may be controlled with Tordon 22K at 2 pt/A sprayed at flowering or in fall.

have vegetative buds that give rise to new shoots. Patch expansion can be dramatic. In Colorado, Dalmatian toadflax shoot density increased over 1,200 percent in six years at one location and 190 percent over three years at another. Yellow toadflax seedlings produce vegetative shoots from root buds two to three weeks after germination. Mature toadflax have well-developed and extensive root systems. Dalmatian toadflax roots may penetrate the soil 4 feet to 10 feet and lateral roots may extend 10 feet from the parent plant; while yellow toadflax roots grow 3 feet deep or more with lateral roots that may extend several yards.

Flowering

Dalmatian toadflax typically flowers beginning in late May or June in Colorado and may continue until fall, particularly if moisture is not limiting. Yellow toadflax begins to flower when shoots are from 16 to 24 inches tall, mid- to late May along the Front Range in Colorado, although at higher elevations (9,000 feet or more), flowering may not begin until late July. Yellow toadflax may not flower until fall under drought conditions. Yellow toadflax shoot phenology in any given patch may range from vegetative to flowering to seed set, depending on the time of season and environmental conditions (particularly moisture). This contributes to management difficulties.

MANAGEMENT

All toadflax species are very difficult to control and management plans should integrate as many strategies as possible to increase potential for success. Assess the condition and composition of the existing plant community in an infested area then determine the approximate composition of the desired plant community needed to achieve land management goals and objectives. Create a management plan that combines various control strategies to foster development of the desired plant community.

Yellow toadflax appears to be more difficult to manage than Dalmatian toadflax. In Colorado, control from Tordon applied at flowering has been most consistent and typically, 4 pt/A is recommended.

Chemical and Cultural Management of Dalmatian Toadflax

Dalmatian toadflax may be controlled with Tordon 22K at 2 pt/A sprayed at flowering or in fall. In Colorado, rates of 2, 4, and 8 pt/A of Tordon were compared and control longevity was greatest from the 2 pt rate, apparently because competition from crested wheatgrass was maintained. Researchers in Wyoming treated Dalmatian toadflax in early September, 1994, with Tordon at 2 pt/A, then seeded the following year in April or August with 'Hycrest' crested wheatgrass, 'Luna' pubescent wheatgrass, 'Critana' thickspike wheatgrass, 'Bozoisky' Russian wildrye, or 'Sodar' streambank wheatgrass. The combination of spraying and seeding competitive grasses controlled Dalmatian toadflax better than spraying alone. Three years after treatments were started, control of Dalmatian toadflax ranged from 61 percent to 86 percent where grasses were seeded in April and from 76 percent to 95 percent from the August seeding, compared to no control from spraying alone.

Chemical Control of Yellow Toadflax

Yellow toadflax appears to be more difficult to manage than Dalmatian toadflax. In Colorado, control from Tordon applied at flowering has been most consistent and typically, 4 pt/A is recommended. Yellow toadflax usually recovers from a single application. For example, Tordon applied at 4 or 8 pt/A controlled 13 percent and 69 percent of yellow toadflax three years after treatments were applied. Other research conducted in Colorado suggests that yellow toadflax control may be improved if Tordon is applied over three consecutive years, but control varied with location. In one experiment conducted at high altitude (Camp Hale; elevation approximately 10,000 feet), 4 pt/A of

Tordon applied at flowering for three consecutive years decreased shoot density to zero. However, the same treatment applied for three years at two other locations (White River drainage, elevation approximately 8,500 feet) controlled 69 percent and 35 percent of yellow toadflax. Telar also may be used to control yellow toadflax in non-crop areas. In an experiment conducted in Middle Park near Parshall, Telar at 1.25 ounce/A applied during flowering or in fall controlled 84 percent of yellow toadflax one year later.

Several classical biocontrol agents are available to use against toadflaxes. However, the success of these agents remains largely unknown.

Escort, 2,4-D amine, Banvel, and Paramount controlled from 5 percent to 24 percent of yellow toadflax one year after single treatments were applied at flowering. Plateau showed some potential to control yellow toadflax in another Colorado experiment where 8 fluid ounces per acre applied once in fall controlled 59 percent of yellow toadflax one year later. While this level of control is unsatisfactory, sequential treatments may increase control but experiments must be conducted to test this hypothesis.

Mechanical and Chemical Control of Yellow Toadflax

Mowing combined with spraying Tordon did not improve control in an experiment conducted near Hesperus, Colorado. Yellow toadflax was mowed three times per year then treated with Tordon at 4 pt/A in fall for two consecutive years and compared to Tordon applied at 4 pt/A at flowering also for two consecutive years. Yellow toadflax control was the same (85 percent) whether Tordon treatments were combined with mowing or not.

Biological Control of Toadflaxes

Several classical biocontrol agents are available to use against toadflaxes. However, the success of these agents remains largely unknown. A defoliating moth (*Calophasia lunula*), an ovary-feeding beetle (*Brachypterosus pulicarius*), and two-seed capsule-feeding weevils (*Gymnaetron antirrhini* and *G.netum*) have been released in the U.S. and Canada to control all toadflax species. Particularly the flowering and seed feeding insects should help decrease seed production.

A stem-boring weevil (*Mecinus janthinus*) and a root-boring moth (*Eteobalea intermediella*) also were released in Canada and the U.S. to control all species of toadflax. These species may help to control shoots and seed production as well as decrease root vigor, but data are unavailable to document their effects. Several of these classical biocontrol agents are available from the Colorado Department of Agriculture Insectary in Palisade. Very few published studies are available to determine whether grazing by livestock will effect any control of Dalmatian or yellow toadflax.

¹ K.G. Beck, Colorado State University Cooperative Extension weed science specialist and professor of weed science, bioagricultural sciences and pest management.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Milan A. Rewerts, Director of Cooperative Extension, Colorado State University, Fort Collins, Colorado. Cooperative Extension programs are available to all without discrimination. No endorsement of products is intended nor is criticism implied of products not mentioned.