



Colorado MASTER GARDENER

Cover Crops and Green Manure Crops no. 7.744

by D. Whiting, C. Wilson, and A. Card'

Outline. . .

The Difference Between a Green Manure and a Cover Crop, page 1

Why is a Cover Crop Beneficial, page 1

Why is Green Manuring Beneficial, page 1

Basic Recipes for the Garden, page 2

Spring Planted, page 2

Fall Planted for Spring Till, page 2

Landscape Uses, page 3

Species Selection, page 3

Annual Species Options, page 3

Perennial Species Options, page 3

Native Species Options, page 4

Establishment and Care, page 4



Putting Knowledge to Work

The Difference Between a Green Manure and a Cover Crop

A **cover crop** is numerous numbers of plants, usually specific annual, biennial, or perennial grasses or legumes, growing and covering the soil surface.

When the cover crop is tilled into the soil it is referred to as a **green manure** crop.

These two terms are often used interchangeably.

Why is Cover Cropping Beneficial?

Cover crops can protect the soil from wind and water erosion, suppress weeds, fix atmospheric nitrogen, build soil structure, and reduce insect pests.

Erosion protection – The primary erosive force in Colorado is wind. Winter winds are especially destructive, carrying away small surface particles of topsoil. A thick stand of a cover crop protects the soil surface from wind erosion and its roots can hold soil in place against water erosion during heavy downpours.

Weed suppression – Cover crops left in place for part or all of a growing season can suppress annual and some perennial weeds. Among the grasses, annual rye has alleopathic properties that prevent weed seeds from germinating and suppress weed seedlings around the root zone of the rye.

Nitrogen fixation – Legumes, inoculated with their specific rhizobium bacteria, will take nitrogen out of the air (present in the soil) and store it in their plant tissues via nodules on the roots of the legume. Some of this nitrogen is available as roots die, but the majority becomes available when the legume is tilled under (green manure).

Soil structure creation – Plant roots exude a sticky substance then glues soil particles together, creating structure. Grasses are exceptional in their ability to do this.

Insect pests reduction – Cover crops encourage beneficial insect populations, often minimizing or eliminating the need for other insect control measures.

Why is Green Manuring Beneficial?

Green manuring enhances soil fertility and soil structure by feeding soil organisms and gluing together soil particles into aggregates.

Soil fertility – When fresh plant material breaks down in the soil, its carbon to nitrogen ratio is low, allowing the nitrogen to be easily released into the soil chemistry by bacteria. Nitrogen accumulated is greater in legumes with rhizobium growing on roots (forming pink nodules), as shown in the following table. Notice the lower figure for rye.

Table 1. Nitrogen accrument of selected cover crops.

Cover Crop	Nitrogen Accrument*
Hairy vetch	3.2 lbs/1000 ft ²
Crimson clover	2.6 lbs/1000 ft ²
Austrian winter pea	3.3 lbs/1000 ft ²
Winter (annual) rye	2.0 lbs/1000 ft ²

* Nitrogen accumulated in growing crop prior to tilling under. Source: ATTRA: Overview of Cover Crops and Green Manures

The following table shows values of nitrogen fixation for legumes. Rates vary due to activity of rhizobium.

Table 2. Potential nitrogen fixation rates of selected legumes for Colorado.

Legume Crop	Pounds N per 1000 ft ²
Crimson clover	1.6-3.0
Field peas	2.0-3.4
Hairy vetch	2.0-4.6
Medics	1.1-2.8
Red clover	1.6-3.4
Sweet clover	2.0-3.9
White clover	1.8-4.6

Source: *Managing Cover Crops Profitably*, Sustainable Agriculture Network

Soil structure – Microorganisms decomposing plant material and the plant material itself produce substances that glue soil particles together. These include slime, mucus and mycelia with the gums, waxes, and resins that aggregate soil particles enhancing the tilth, porosity, and water holding capabilities of soil.

Basic Recipes for Cover Crops and Green Manure Crops in a Garden

Spring Planted

Most gardeners don't have enough space to forfeit to a cover crop for an entire growing season. But if you do a spring seeded clover it would give your soil a great boost. Some seed companies will *rhizo-coat* seed with the specific rhizobium for you. If not, apply rhizobium as specified on the bag. Rhizobium comes in a black powder specific to the species of clover. It also has a definite shelf life, so check the expiration date. Broadcast the rhizobium mix at a specified rate after the last frost with a hand held broadcaster (often used with pelleted fertilizer) into a loose seedbed and incorporate shallowly and water until germinated. Monitor water as you would in a lawn.

Till under at least two weeks prior to planting. Decomposing plant material consumes soil oxygen and can create plant health problems if not tilled in ahead of time. More than one tilling may be necessary to get an acceptable kill of the clover.

Fall Planted for Spring Till

Most will opt for a fall cover crop tilled under as a spring green manure. Seeding dates should be by mid-October at the latest. Mid-September is ideal on the Front Range. In mountain elevations, plant in August or earlier. A rye/ Austrian winter pea or rye/hairy vetch mixture will overwinter in Colorado. Hairy vetch is hardier than winter pea. Rye is extremely winter hardy. Prepare as above and broadcast at the following specified rates.

Table 3. Seeding rates for selected winter cover crops.

Cover Crop	Ounces per 100 square feet	Pounds per 1000 square feet
Winter Rye	4-6	2.5-3.75
Austrian Winter pea	4-6	2-4
Hairy Vetch	2-3	1-2

Source: *Managing Cover Crops Profitably*, Sustainable Agriculture Network

Overwintered cover crops become a veritable salad bar to geese and deer. A cover crop that is well established prior to winter temperature extremes should rebound from wildlife grazing in late winter or early spring.

Till the cover crop in mechanically or turn it under with a spade a month before you plan to plant/seed into that area. Decomposing plant material consumes soil oxygen and can create plant health problems if not tilled in ahead of time.

Landscape Uses

Bare soil presents erosion and aesthetic issues for homeowners. During droughty periods watering restrictions and natural precipitation may make turf establishment difficult or impossible. A temporary cover crop or long term xeric grass may be the answer.

In this scenario, the homeowner should understand that a cover crop will not look or feel like a healthy Kentucky bluegrass lawn, but should satisfy the need to cover the soil.

Species Selection

Annual Species Options

These are cool season grains that should be broadcast at 2 to 3 pounds per 1000 square feet in February or March. Natural precipitation may be sufficient to get them established. They are suited for non-traffic areas, as they will grow to 2 feet tall and brown-out in the heat of summer. The Sterile Triticale will not produce viable seeds so may be a good idea for areas that will eventually be put into turf or garden space. Winter rye seeds can be a weed problem in seeded turf grass and gardens.

Table 4. Types of annual grasses.

Name	Bunch or Sod	Cool or Warm Season	Annual or Perennial	Turf?	Reseed?
Winter rye	Bunch	Cool	Annual	No	Yes
Pioneer Sterile Triticale	Bunch	Cool	Annual	No	No

Perennial Species Options

These are non-native grasses often used on roadsides for stabilization and cover. They are perennial and will be persistent (i.e., difficult to kill) once they are established. Water requirements for both are 9 to 10 inches of precipitation per year. Streambank wheatgrass has a slightly higher water requirement but is tolerant of very clayey soils, unlike Crested wheatgrass. Broadcast in February or March at 3 to 5 pounds per 1000 square feet.

Table 5. Types of perennial grasses.

Name	Bunch or Sod	Cool or Warm Season	Annual or Perennial	Turf?	Reseed?
Streambank Wheatgrass	Sod	Cool	Perennial	Yes	Some
Crested Wheatgrass	Bunch	Cool	Perennial	Yes	Some

Native Species Options

These have the lowest water requirements at 8 inches of precipitation per year and should be considered for areas of a landscape that are being converted to xeric management. This is a longterm management decision as the price of these seeds is more than the other options. These will not feel like Kentucky Blue grass and will brown-out like other cool season grasses. Seed as per perennial species options specifications. Seed for native species will be available from local seed sources, such as Pawnee Buttes Seeds.

Table 6. Types of native grasses.

Name	Bunch or Sod	Cool or Warm Season	Annual or Perennial	Turf?	Reseed?
Indian Ricegrass	Bunch	Cool	Perennial	No	Some
Squirreltail Bottlebrush	Bunch	Cool	Perennial	No	Some

Establishment and Care

Before seeding – Prepare a seedbed for fine grass seed, ideally amending the soil with compost and tilling as deeply as possible. If possible, fence off the area from traffic.

Seeding – Water area prior to seeding if possible to establish ample soil moisture levels.

Broadcast the correct amount of seed per area onto a loosely tilled, fine (no soil pieces bigger than ¼ inch) seedbed. Shallowly incorporate seed with garden rake (not a leaf rake) to a depth of ¼ to ¾ inch deep.

For larger areas consider hydromulching the seed. This will save time and increase germination of seeds.

After seeding – Consider laying a thin layer (less than 1 inch deep) of seed free straw to hold in moisture and increase germination and survival of grass seedlings. Bird netting over the straw fastened to the ground with landscape fabric staples will keep the straw from blowing away.

Check moisture levels in the upper inch of soil at least every other day (soil should feel as moist as a wrung out sponge) and water if necessary (and if possible).

Mowing – If necessary, mow as high as possible.

Removing cover crops – For Winter Rye, either till under, mow and mulch heavily, or spray herbicide before it goes to seed. A seed bank can be sodded over or watered, germinated and killed. Perennial grasses can be either mowed and mulched heavily prior to sodding or sprayed with herbicide and sodded or sprayed with herbicide, tilled and seeded.

Colorado Master Gardener training is made possible, in part, by a grant from the Colorado Garden Show, Inc.

¹D. Whiting, Colorado State University, Cooperative Extension consumer horticulture specialist and Colorado Master Gardener coordinator; C. Wilson, Extension horticulture agent, Denver County; and A. Card, Extension horticulture agent, Boulder County.

Colorado State University, U.S. Department of Agriculture and Colorado counties cooperating. Cooperative Extension programs are available to all without discrimination. No endorsement of products mentioned is intended nor is criticism implied of products not mentioned.