



# RANGE

---

## Diffuse and Spotted Knapweed

no. 3.110

by K.G. Beck <sup>1</sup>

### Quick Facts...

Diffuse knapweed is a short-lived, non creeping perennial, a biennial, or occasionally an annual that reproduces and spreads solely from seed.

Spotted knapweed is a short-lived, noncreeping perennial that reproduces from seed and forms a new shoot each year from a taproot.

Diffuse and spotted knapweed are readily controlled with herbicides. Unless cultural techniques are used, however, the weeds will reinvade.

Diffuse knapweed (*Centaurea diffusa*) is a short-lived perennial, a biennial, or occasionally an annual. It reproduces and spreads from seed. The plant develops a single shoot (stem), 1 to 2 feet tall, that is branched toward the top. Grazed plants may produce multiple stems. Rosette and lower shoot leaves are finely divided. Leaves become smaller toward the top of the shoot and have smooth margins.

Many solitary flowering heads occur on shoot tips. They are about 1/8 inch in diameter and 1/2 to 2/3 inch long. Flowers usually are white but may be purplish. Involucre bracts are divided like teeth on a comb and tipped with a slender spine that makes them sharp to the touch. Sometimes the bracts are dark-tipped or spotted like spotted knapweed. The long terminal spine differentiates diffuse from spotted knapweed.

Spotted knapweed (*Centaurea maculosa*) looks like diffuse knapweed with some notable exceptions. Spotted knapweed is a short-lived, noncreeping perennial that reproduces from seed (primary means of spread) and forms a new shoot each year from a taproot. The weed produces one or more shoots that are branched and 1 to 3 feet tall. Rosette leaves can be 6 inches long and deeply lobed. Leaves are similar to diffuse knapweed. Lavender to purple flowers are solitary on shoot tips and about the same size as diffuse knapweed flowers. Involucre bracts are stiff and black-tipped. The tip and upper bract margin have a soft, spine-like fringe and the center spine is shorter than others.

### Phenology, Biology and Occurrence

Diffuse knapweed seeds germinate in spring or fall or anytime during the growing season following a disturbance, if adequate soil moisture is present. Seedlings develop into rosettes and diffuse knapweed remains as a rosette until it grows to a critical size, then it bolts, flowers, and sets seed. It may take from one to several years for diffuse knapweed to reach the critical size necessary to reproduce by seed.


**Diffuse knapweed** is native to degraded noncropland (waste places) and seashores from southern Europe to north-central Ukraine. It generally is found on dry, light, porous soils in Europe. Diffuse knapweed appears to occupy similar areas in the United States. Diffuse knapweed will not tolerate flooding or shade and thrives in the semiarid west (generally in 9- to 16-inch precipitation zones). Environmental disturbance (e.g., overgrazed pastures or rangeland, roadsides, rights-of-way, gravel piles, etc.) promotes its invasion.

In Colorado, the worst infestations occur along the Front Range in Larimer, Boulder, Douglas and El Paso counties. Severe infestations also occur in Archuleta and La Plata counties. A 2002 survey conducted by the Colorado Department of Agriculture found 145,148 infested acres of diffuse knapweed and 1,093 infested acres of spotted knapweed.

**Colorado  
State**  
University  
Cooperative  
Extension

*Putting Knowledge to Work*

© Colorado State University  
Cooperative Extension. 3/03.  
[www.ext.colostate.edu](http://www.ext.colostate.edu)



**Spotted knapweed** germinates in spring or fall. Perennial plants resume growth in early spring and bolt at approximately the same time as diffuse knapweed. Flowering occurs through the summer into fall.

Spotted knapweed is native to central Europe, where it is found in light, porous, fertile, well-drained and often calcareous soils in warm areas. It occupies dry meadows, pastureland, stony hills, roadsides, and the sandy or gravelly floodplains of streams and rivers. Spotted knapweed tolerates dry conditions, similar to diffuse knapweed, but survives in higher moisture areas as well (e.g., it thrives in the wetter conditions of the western Montana mountains). Spotted and diffuse knapweed infestations often occur together in Colorado.

## Management

Diffuse and spotted knapweed can be managed similarly. They are readily controlled with herbicides. However, the weeds will reinvade unless cultural techniques are used.

**Chemical control.** Research conducted at Colorado State University indicates that Tordon 22K (picloram) at 1 to 2 pt/A, Transline (clopyralid) at 0.67 to 1 pt/A, Curtail (clopyralid + 2,4-D) at 4 to 6 pt/A, or Banvel/Vanquish/Clarity (dicamba) at 1 to 2 pt/A control diffuse knapweed. Tank mixes of Banvel/Vanquish/Clarity plus 2,4-D at 1 pt + 2 pt/A or Banvel/Vanquish/Clarity plus Tordon 22K at 1 to 2 pt + 0.5 to 1 pt/A or Tordon plus 2,4-D at 0.75 pt + 2 pt/A all control diffuse knapweed. These tank-mixes may save money and reduce grass injury resulting from higher use rates of a single herbicide.

Spotted knapweed and diffuse knapweed generally occupy the same areas in Colorado, so the same herbicide treatments can be applied. Weed scientists at Montana State University indicate that 1 pt/A of Tordon (0.25 lb) controls spotted knapweed for two to three years, but the weed will reinvade the area unless other management techniques are used.

**Cultural control.** If desirable grass competition is evident in diffuse or spotted knapweed stands, judicious herbicide application that does not injure grasses may allow them to compete effectively with the weeds. Irrigation (where possible) may help stimulate grass competition in these cases. However, infested rangeland or pastures often are degraded, allowing knapweed invasion, and herbicides alone will not restore the land to a productive state. Seeding suitable perennial grasses is necessary to prevent weed reinvansion.

**Biological control.** Many insects are being evaluated for biological control of diffuse and spotted knapweeds. Researchers at Montana State University believe it will take a complex of insects (perhaps 12) to reduce diffuse and spotted knapweed populations.

Several insects are available in Colorado, from the Colorado Department of Agriculture. The seedhead flies *Urophora affinis* and *U. quadrifasciata* have been released in many Front Range counties. These insects cause plants to produce fewer viable seeds and abort terminal or lateral flowers.

Root-feeding insects may have a more detrimental effect on knapweed populations than seed-feeding ones. Larvae of the diffuse knapweed root beetle (*Sphenoptera jugoslavica*) feed in the roots of diffuse knapweed. Larvae of the yellow-winged knapweed moth (*Agapeta zoegana*) feed and the knapweed root weevil (*Cyphocleonus achates*) in the roots of both knapweed species.

Spotted knapweed infestations are not as severe in Colorado as diffuse knapweed. However, this weed spreads rapidly. For example, spotted knapweed was first observed in Gallatin County, Montana, in the 1920s, but is now found in all Montana counties. Today, over 4.7 million acres are infested.

Livestock (sheep, goats, cattle) will eat diffuse and spotted knapweed. Recent research completed by Colorado State University shows that cattle grazing diffuse knapweed twice in spring decreased seed set by 50 percent and tumbling off-site over winter by 15 percent. Cattle were managed to achieve 50 percent utilization of pasture and were allowed to graze at two 10-day intervals when diffuse knapweed was bolting and about 6 to 12 inches tall.

<sup>1</sup> Colorado State University Cooperative Extension weed specialist and professor, plant pathology and weed science.