

Dalmatian toadflax

Colorado Dept. of
Agriculture,
Conservation
Services Division
700 Kipling Street
Suite 4000
Lakewood, CO 80215
303-239-4100



Key ID Points

1. Showy yellow snapdragon-like flowers with an orange throat on elongated racemes.
2. Thick, waxy, bluish heart-shaped leaves that wrap the stem.

Dalmatian toadflax Identification and Management



Identification and Impacts


Dalmatian toadflax (*Linaria dalmatica*) is a non-native, perennial forb introduced from the Mediterranean region as a folk remedy, fabric dye and ornamental. It reproduces both by seed and by extensive, creeping rhizomes. A single plant produces 500,000 seeds, most of which fall within 18 inches of the parent plant. Seeds can remain viable for at least 10 years. Dalmatian toadflax grows to 3 feet, and has bright yellow snapdragon-like flowers with an orange throat on elongated racemes. The alternate leaves are broad, with a thick, waxy cuticle and a bluish cast. Each leaf is heart-shaped and wraps the stem.

Habitats for Dalmatian toadflax include disturbed open sites, fields, pastures, rangeland, roadsides, cropland and forest clearings. Infestations can begin in small disturbed sites, then spread even to rangeland and wildlife habitats in excellent condition. Dalmatian toadflax is a highly aggressive plant that can genetically adapt to varied environmental conditions and herbicide controls. Its extreme competitiveness is due to early spring regeneration from vegetative buds on roots that are not dependent on soil moisture or native plant competition. Once established, toadflax quickly overruns native plants and becomes

a monoculture that severely reduces forage, productivity, biodiversity and wildlife habitat.

The key to effective control of Dalmatian toadflax is prevention and integrating as many management strategies as possible. Prevention is always desirable when dealing with Dalmatian toadflax. Early detection and eradication can keep populations from exploding, making more management options available. With the plants varying genetically using many different approaches is important such as; chemical, mechanical, cultural and biological methods. Details on the back of this sheet can help to create a management plan compatible with your site ecology.

Dalmatian toadflax is designated as a "List B" species in the Colorado Noxious Weed Act. It is required to be either eradicated, contained, or suppressed depending on the local infestations. For more information visit www.colorado.gov/ag/csd and click on the Noxious Weed Management Program. Or call the State Weed Coordinator at the Colorado Department of Agriculture, Conservation Services Division, 303-239-4100.



Clockwise, from lower left, photos © John M. Randall of The Nature Conservancy; and Linda Wilson and Susan Turner of Invasive.org. Infestation map by Crystal Andrews, Colorado Department of Agriculture.

Linaria dalmatica

**CULTURAL**

It is imperative to seed managed areas with competitive grasses such as thickspike wheatgrass and streambank wheatgrass. The combination of herbicide spraying and seeding competitive grasses controls Dalmatian toadflax better than spraying alone. (K.G. Beck, CSU)

**BIOLOGICAL**

Calophasia lunula, a predatory noctuid moth, feeds on leaves and flowers of Dalmatian toadflax. *Eteobalea intermediella*, a root boring moth, and *Mecinus janthinus*, a stem boring weevil, are also available. For more information, contact the Palisade Insectary of the Colorado Department of Agriculture at 970-464-7916.

**MECHANICAL**

For small infestations, pulling toadflax by hand can be effective. Pull every year for 5 to 6 years to deplete the reserves of the root system. Monitor the site for 10 - 15 years to remove seedlings produced from dormant seeds.

Integrated Weed Management:

Because of the high genetic variability of the toadflax species, it is critical to integrate as many management strategies as possible into the control program. Two local populations may respond differently to the same herbicides.

Keys to management are to prevent seed formation and vegetative spread by roots.

Controlling toadflaxes is expensive and difficult, prevention is the best option.

HERBICIDES

NOTE: The following are recommendations for herbicides that can be applied to range and pasturelands. Rates are approximate and based on equipment with an output of 30 gal/acre. Please read label for exact rates. **Always read, understand, and follow the label directions. The herbicide label is the LAW!**

HERBICIDE	RATE	APPLICATION TIMING
Picloram (Tordon 22K) *Avoid spraying near trees and/or water	2-4 pints/acre	Apply when flowering and/or in the fall. Add non-ionic surfactant @ 0.32oz/gal water or 1 qt/100 gal water.
Chlorsulfuron (Telar)	2-3 oz./acre	Apply at flower stage in spring and/or in the fall. Add non-ionic surfactant @ 0.32 oz/gal. water or 1 qt./100 gal. water.
2,4-D + Dicamba	2 qt. + 2 qt/ acre	Apply during pre-bloom to flower stage. Add non-ionic surfactant @ 0.32 oz/gal. water or 1 qt/100gal water. Retreatment is required for several years.

Top photo, © Kelly Uhing, Colorado Department of Agriculture. *Calophasia lunula* larva photo © Bob Richard, USDA APHIS, Invasive.org. Handpulling toadflax photo © Lake Tahoe Environmental Education Coalition.

Dalmatian toadflax

