

UCS 20/622/8010

SERVICE c.2

RECEIVED

IN ACTION

MAY 24 1990

COLORADO STATE LIBRARY
State Publications Library

COLORADO STATE UNIVERSITY EXTENSION SERVICE

Edema disease in swine

Martin Bergeland and
Harold Kurtz^{1/}

no. 8.010



Quick Facts

- Edema disease is characterized by an excessive accumulation of fluid in body tissues.
- Edema usually afflicts thrifty and fast-growing pigs and usually strikes shortly after weaning.
- Various outward symptoms, such as staggering gait, head tilting, odd lying and sitting positions and other signs indicate the disease's presence.
- Edema results from an increase in certain *E. coli* bacteria in the animal's small intestine shortly after weaning.
- Accurate diagnosis is important since other diseases produce similar symptoms.
- Diet changes can be employed as treatment and control.

Symptoms

One or more pigs may be found dead unexpectedly as a result of the disease. At the same time, one usually finds pigs with nervous symptoms, including staggering, head tilting, stumbling and falling, assuming a "dog sitting" position, lying on the sternum, or lying on the side and making continuous kicking movements.

Affected pigs do not have a fever unless the environmental temperature is unusually high. Some pigs may have swollen eyelids. The mortality rate among pigs with symptoms is about 65 percent. Those that do not die show improvement after two days.

The course of the disease is about two to five days, but in some herds, there is a recurrence 10 days to two weeks later. In this second outbreak, affected pigs often walk slowly or wander aimlessly around the edge of the pen, with the head slightly elevated or tilted to one side.

Cause

The disease symptoms result from rapid growth of certain types of bacteria, *Escherichia coli*, in the small intestine. After weaning, the number of *E. Coli* in the intestine tends to increase, usually reaching a maximum in about four days post-weaning. There are many different types of *E. coli* most of which do not cause edema disease. But if there are edema disease-producing strains of *E. coli* in the intestine, these also will increase during the post-weaning period and may become the predominant bacteria in the small intestine.

Edema disease-producing strains of *E. coli* produce one or more toxins that are absorbed from the intestine into the blood. These toxins damage blood vessels and affect blood pressure, which in turn causes fluid to leak from vessels and accumulate in many body tissues. This fluid accumulation is most important in the brain,

Edema disease usually occurs three days to two weeks after weaning, but it is seen occasionally in nursing pigs. Affected pigs develop nervous symptoms or die suddenly.

Edema is an excessive accumulation of fluid in body tissues. The disease was named "edema disease" or "gut edema" because those who first observed the disease in the 1930s found excess fluid in the walls of the stomach and intestine or under the eyelids of some pigs with this problem. Pigs with edema disease may have excess fluid in many parts of the body, but involvement of the brain is most important and causes the clinical symptoms.

Incidence

Edema disease can affect pigs 3-14 weeks old but usually occurs within three days to two weeks after weaning. The affected pigs often are the most thrifty and fast-growing individuals of a group. Edema is commonly seen under conditions of excellent management and nutrition. The morbidity rate usually is about 15 percent, but 50 percent or more of some groups are affected.

^{1/} Martin Bergeland, South Dakota State University, and Harold Kurtz, University of Minnesota. This information, originally published in the **Pork Industry Handbook**, has been reviewed for Colorado producers by Marvin Heeney, CSU extension professor, animal sciences (4/15/79)

where swelling can result in destruction of some brain tissue and, in many cases, death of the animal.

Differential Diagnosis

There are many diseases of young pigs that produce symptoms similar to edema disease. Some of the more common problems include the following:

- Virus infections (pseudorabies, hemagglutinating encephalomyelitis virus, polioencephalomyelitis)
- Water deprivation (salt poisoning)
- Bacterial infections (Glasser's disease, bacterial meningitis, ear infection)
- Porcine stress syndrome
- Mulberry heart disease
- Chemical poisons (arsenic, lead, mercury, various insecticides and rodenticides)

It is important to obtain an accurate diagnosis, since treatment and preventive measures are specific for each of these different diseases.

Treatment and Control

An abrupt change in the ration may help to prevent additional cases. Changing the diet will alter the growth conditions for bacteria in the

intestine, and it may allow other types of bacteria to proliferate and replace the strains of *E. coli* that cause edema disease. (Even though a change in the diet may help to stop the outbreak, this does not imply that the original ration was improperly formulated. Changing the feed simply is an effort to alter the conditions for bacterial growth within the intestine.)

Withholding feed temporarily or feeding soaked whole oats for one to two days are methods that have been used to change the diet suddenly. The original ration may be restored gradually over a period of four to five days.

Treatment with antibacterial drugs also may help prevent additional cases. Treatment of pigs that have clinical symptoms, however, often is ineffective.

In some herds, edema disease tends to occur in each successive group of pigs that is weaned. The use of antibiotics or sulfonamides in the feed or water at weaning time may be beneficial in these herds.

Vaccines produced from edema disease-producing strains of *E. coli* generally have been ineffective in preventing the problem.

Treatments or changes in management practices should be designed to meet the needs of each individual production unit. Consult a veterinarian to obtain recommendations for individual herds.