



Immunization Quick Reference Chart for Child Care, Preschool & K-Entry



1. Select the appropriate age range for the child from the left hand column. Review the child's immunization record to assure they have at least the number of doses listed under each column directly to the right of their age.
2. If the child does not have the minimum number of doses, they must immediately be placed on a plan to obtain the remaining vaccinations. The parent should be directly notified (in person or by phone or mail) that their child does not have the minimum number of required doses. The parent must complete and return a plan for immunizations within 14 days of the direct notification. (Remember: Colorado law has allowances for medical, religious and personal exemptions).
3. Share this chart with parents, other childcare providers and licensing staff so they can easily monitor children's immunization needs.

Age of Child	# of required doses DT, DTP, or DTaP <i>Diphtheria, Tetanus Pertussis</i>	# of required doses Polio <i>Polio</i>	# of required doses MMR <i>Measles Mumps Rubella</i>	# of required doses Hib <i>Haemophilus influenzae type b</i>	# of required doses Hep B <i>Hepatitis B</i>	# of required doses Varicella <i>Chickenpox</i>	# of required doses PCV7 <i>Pneumococcal Disease</i>
< 4 mos.	0	0	0	0	0	0	0
by 4 mos.	1	1	0	1	1	0	1
by 6 mos.	2	2	0	2	2	0	2
by 8 mos.	3	3	0	2	2	0	3/2•
by 12 mos.	3	3	0	3/2♥	2	0	3/2•
by 15 mos.	3	3	1	3/2♥	3	1	4/3/2•
by 18 mos.	4	3	1	3/2/1♥	3	1	4/3/2•
by 2 years	4	3	1	3/2/1♥	3	1	4/3/2•
by 3 years	4	3	1	3/2/1♥	3	1	0
by 4 years	4	3	1	3/2/1♥	3	1	0
K-Entry	5/4♦	4/3♣	2	0	3	2	0

- ♥ The number of Hib doses required depends on the child's current age and the age when the Hib vaccine was administered. If any dose is given ≥ 15 months, the Hib requirement is met. For students who begin the series < 12 months, 3 doses are required, of which at least 1 dose must be administered at ≥ 12 months. If the 1st dose was given at 12 to 14 months, 2 doses are required. If the current age is ≥ 5 years, no new or additional doses are required.
- The number of pneumococcal conjugate vaccine (PCV7) depends on the student's current age and the age when the 1st dose was administered. If the 1st dose was administered between 2 to 6 months of age, the child will receive 3 doses two months apart and an additional dose between 12 to 15 months of age. If started between 7 to 11 months of age, the child will receive 2 doses two months apart and an additional dose between 12 to 15 months of age. If the series was started between 12 to 23 months of age, then the child will receive 2 doses, two months apart. There is no requirement once the child turns two years of age.
- ♦ If the 4th dose of DTaP was given ≥ 4 years, only 4 doses are required.
- ♣ If the 3rd dose of Polio was given ≥ 4 years, only 3 doses are required.